

OPEN HERE

READ THE LABEL BEFORE USING
(SOMA MAELEZO KABLA YA KUTUMIA)

A pre or post-emergence herbicide with some soil residual activity for broad-leaved weeds control and some grasses in wheat and barley.

(Dawa ya kuua kwekwe kabla ya mimea kuota au baada ya kuota, ambayo huangamiza kwekwe wenye majani mapana na baadhi ya nyasi kwenye ngano na shayiri).

COMMERCIAL AND AGRICULTURAL CLASS**GUARANTEE (DHAMANA):**

Chlorsulfuron: 75%
Inert ingredients 25%

REGISTRATION NO.: **PCPB (CR) 0073**
(NAMBARI YA USAJILI)

Shelf Life: 3 years from the date of manufacture if stored in unopened original container in a dry cool place

Maisha rafuni: Miaka mitatu baada ya tarehe ya kutengenezwa iwapo dawa hii itahifadhiwa kwa chombo chake asili bila kufunguliwa na iwe mahali pakavu na pasipo na joto.



CAUTION
ILANI



Batch No.:
(Nambari ya furushi)
Date of Manufacture:
(Tarehe ya kutengenezwa)

In case of emergency call toll free numbers / Wakati wa dharura piga simu bila malipo 0800 730030/ 0800 720021 (24hrs/masaa 24)

Registrant / Wasajiliwa:

DuPont International Operations Sàrl (DIOSàrl)
2, Chemin du Pavillon - CH-1218 Le Grand-Saconnex,
Geneva, Switzerland

Manufacturer / Watengenezaji:

Du Pont de Nemours (FRANCE) S.A.S
82 rue de Wittelsheim - 68701 Cernay - France

Agent: Du Pont De Nemours Kenya Limited

P.O Box 29171- 00100, Nairobi, Kenya.
Tel: + 254 20367500

Distributor / Msanzaji: Elgon Kenya Limited

Eastgate Rd. Off. Mombasa Rd. - P.O.Box 46826
00100 - Nairobi (Kenya)
Tel: +254-20-6534410
Fax: +254-20-6534807
www.elgonkenya.com



® Registered trademark of (Alama ya biashara ya kampuni ya) Du Pont de Nemours & Co. (Inc).



DU PONT

Glean® 75 DF

HERBICIDE / CHLORSULFURON

GROUP

B

HERBICIDE

KEEP LOCKED OUT OF REACH OF CHILDREN
(FUNGIA MBALI NA WATOTO)



NET CONTENTS (Kiasi):
100 g

P 358 **CYAN** **MAGENTA** **JAUNE** **NOIR**

K-35530/31410 - KENYA

**READ THE LABEL BEFORE USING
(SOMA MAELEZO KABLA YA KUTUMIA)**

A pre or post-emergence herbicide with some soil residual activity for broad-leaved weeds control and some grasses in wheat and barley.

(Dawa ya kuua kwekwe kabla ya mimea kuota au baada ya kuota, ambayo huangamiza kwekwe wenye majani mapana na baadhi ya nyasi kwenye ngano na shayiri).

COMMERCIAL AND AGRICULTURAL CLASS

GUARANTEE (DHAMANA):

Chlorsulfuron: 75%
Inert ingredients 25%

REGISTRATION NO.: **PCPB (CR) 0073**
(NAMBARI YA USAJILI)

Shelf Life: 3 years from the date of manufacture if stored in unopened original container in a dry cool place

Maisha rafuni: Miaka mitatu baada ya tarehe ya kutengenezwa iwapo dawa hii itahifadhiwa kwa chombo chake asili bila kufunguliwa na iwe mahali pakavu na pasipo na joto.

In case of emergency call toll free numbers / Wakati wa dharura piga simu bila malipo 0800 730030/ 0800 720021 (24hrs/masaa 24)

Registrant / Wasajiliwa:

DuPont International Operations Sàrl (DIOSàrl)
2, Chemin du Pavillon - CH-1218 Le Grand-Saconnex,
Geneva, Switzerland

Manufacturer / Watengenezaji:

Du Pont de Nemours (FRANCE) S.A.S
82 rue de Wittelsheim - 68701 Cernay - France

Agent: Du Pont De Nemours Kenya Limited

P.O Box 29171- 00100, Nairobi, Kenya.
Tel: + 254 20367500

Distributor / Msanzaji: Elgon Kenya Limited

Eastgate Rd. Off. Mombasa Rd. - P.O.Box 46826
00100 - Nairobi (Kenya)
Tel: +254-20-6534410
Fax: +254-20-6534807
www.elgonkenya.com



® Registered trademark of (Alama ya biashara ya kampuni ya) Du Pont de Nemours & Co. (Inc).



Glean® 75 DF

HERBICIDE / CHLORSULFURON

GROUP

B

HERBICIDE

**KEEP LOCKED OUT OF REACH OF CHILDREN
(FUNGIA MBALI NA WATOTO)**



**NET CONTENTS (Kiasi):
100 g**




**CAUTION
ILANI**



K-35530/31410 - KENYA

P 358 **CYAN** **MAGENTA** **JAUNE** **NOIR**





In case of poisoning, call the 24 hour Toll Free Lines 0800-720021/ 0800730030 for professional advice.

® Registered trademark of Du Pont de Nemours & Co. (Inc).

GENERAL INFORMATION

GLEAN® 75 DF is recommended on all varieties of wheat and barley. It is active both through root uptake and by contact through the foliage. Weed control is most effective either pre-emergence or when the weeds are small and actively growing. GLEAN® 75 DF can cause severe stunting of more resistant weeds which subsequently do not compete with the crop.

RECOMMENDATIONS FOR USE

GLEAN® 75 DF is recommended at 20 g product per hectare for post-emergence applications and 12.5 - 15 g in tank-mix, applied early post-emergence. Rate for 20L of water (Knapsack sprayer): 4g of product. Spray volume varies between 100- 600L of water per Hectare depending on crop and weed growth stage. Higher weed density requires higher spray volume to ensure proper coverage.

PRE HARVEST INTERVAL: Do not treat later than BBCH 30 (Beginning of stem elongation).

TIMING OF APPLICATION

Barley

For best results GLEAN® 75 DF should be applied post-emergence to barley at 2-3-leaf stage till the end of tillering.

NOIR

Wheat

GLEAN® 75 DF is preferably applied pre-emergence, also early post-emergence, also early post-emergence at 2-3 leaf stage till end of tillering. Avoid overdosing caused by slow movement, turning, stopping and overlapping.

NB: Symptoms appear after 3-10 days, with necrosis and death of the weeds taking 2 to 4 weeks depending on growing conditions.

PREPARATION OF THE SPRAY MIXTURE

Half fill spray tank with water. Premix the required amount of GLEAN® 75 DF with a small amount of water. Add this premix to the tank. Top up tank with water to the required volume. Agitate and mix thoroughly. Do not keep spray mixture for over 12 hours.

CLEANING OF SPRAYING EQUIPMENT

It is important to thoroughly clean all traces of GLEAN® 75 DF from application equipment immediately after use. First drain tank, then flush tank, pump, hoses and boom twice with a solution of 1/2 L chlorine bleach (5% sodium hypochlorite) per 100 L water. Leave this solution during 15 minutes under agitation in sprayer. Remove nozzle tips and filters and clean separately. To terminate, rinse tank, hoses and boom with clean water to remove traces of chlorine bleach.

FOLLOWING CROPS - CAUTION

Do not plant any other crop, except wheat or barley, within 18 months of spraying GLEAN® 75 DF. GLEAN® 75 DF should not be sprayed on cereals that are undersown with pastures.

Crop failure

In the event of crop failure for any reason, replant with wheat or barley only.

WEEDS CONTROLLED

The following weeds are controlled by GLEAN® 75 DF at the recommended rate and timing of application.

Common name

Oxalis
Spurrey
Chickweed
Wild raddish (white charlock)
Black bindweed
Round-leaved Goosefoot
Prostrate knotweed
Shepherd's purse

Scientific name

Oxalis spp
Spergula arvensis
Stellaria media
Raphanus raphanistrum
Polygonum Convolvulus
Chenopodium opulifolium
Polygonum aviculare
Capsella bursa-pastoris

Pigweed
Mallow
Chinese lantern
Mexican marigold
Gallant soldier
Devils thorn
Black knightshade
Thorn apple
Blackjack
Portulaca
Volunteer Rape Double thorn
Rye grass

Amaranthus spp
Malva verticillata
Nicandra physalodes
Tagetes minuta
Galinsoga parviflora
Emex australis
Solanum nigrum
Datura stramonium
Bidens pilosa
Portulaca oleracae
Oxygonum sinuatum
Lolium spp.

PRECAUTIONS

Avoid contact with skin, eyes and clothing. May irritate eyes, nose, throat and skin.

Do not inhale dust or spray mist.

Keep away from children and do not eat, drink of smoke while mixing and applying.

Wear protective clothing, gloves and goggles while preparing spray and spraying.

After work

Wash gloves and hands with soap and plenty of water and change clothing.

STORAGE

Store product in closed original container in a cool dry place and keep away from children and foodstuffs. Keep away from food/feedstuffs, fertilizer, other pesticides and seed.

DISPOSAL

Do not re-use empty container pack for any other purpose. Puncture and burn or bury the container in an authorized place.

ENVIRONMENTAL HAZARDS

Do not dispose of spray mixture near waterways. Avoid spray drift into susceptible crops, grazing, rivers, dams and areas not under treatment. Do not spray, drain or flush equipment on or near desirable trees or plants where their roots may extend.

FIRST AID INSTRUCTIONS

In case of contact with eyes: Flush with water for a minimum of 10 minutes. Seek medical attention.



In case of Contact with skin: Wash with soap and plenty of water. Seek medical attention.

If swallowed: Induce vomiting. Seek medical attention.

If inhaled: Move to fresh air. Seek medical attention.

TOXICOLOGICAL INFORMATION /NOTICE TO PHYSICIAN

Antidote: There is no known antidote

Treatment: In case of poisoning, apply symptomatic treatment.

In case of emergency call toll free numbers 0800 730030/0800 720021 (24hrs)




NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.





Panapo ajali ya kusumishwa, piga nambari za simu zifuatazo zilizowazi kwa masaa 24 na bila malipo kwa ushauri maalum: 0800720021 au 0800730030

® Alama ya biashara ya kampuni ya Du Pont de Nemours & Co. (Inc).

MAELEZO KWA JUMLA

GLEAN® 75 DF inapendekezwa kutumiwa kwa aina zote za ngano na shayiri. Huingia katika mimea kupitia mizizi pamoja na majani yake; kuweza kuzuia kwekwe, inafaa kunyunyizia dawa wakati kwekwe hao bado wachanga. GLEAN® 75 DF pia inaweza kuzuia kukua kwa kwekwe ambao wanabaki baada ya kunyunyizia dawa.

JINSI YA KUTUMIA

GLEAN® 75 DF inapendekezwa kutumika katika vipimo vya gramu 20 kwa kila Hektari au gramu 12.5 hadi 15 ikiwa utachanganya na dawa zinginezo na unyunyizie mapema. Au changanya gramu 4 za GLEAN® 75 DF kwa Lita 20 za maji.

Tumia 100-600L ya maji kwa hektari. Maji mengi hutumika kuchanganya dawa ikiwa idadi ya magugu ni juu ili kuhakikisha dawa imesambazika vyema.

Muda wa kusubiri kabla ya kuvuna

Usinyunyize GLEAN® 75 DF wakati mimea imeendelea kumea na matawi kurefuka. (BBCH 30)

Muda wa kusubiri kabla ya kuvuna

Usinyunyize GLEAN® 75 DF wakati mimea imeendelea kumea na matawi kurefuka. (BBCH 30)

WAKATI WA KUNYUNYIZIA

Shayiri: Kwa matokeo bora, GLEAN® 75 DF inatajikana kwa shayiri kutumika kuua kwekwe baada ya mimea kuota "*post-emergence*" (wakati mimea ina majani 2 au 3 hadi mwisho wa "*tillering*").

Ngano: Kwa ngano GLEAN® 75 DF inatakiwa kutumika kwa matokeo mema zaidi kuua kwekwe kabla ya mimea kuota "*pre-emergence*" au baada ya mimea kuota "*post-emergence*" (wakati mimea ina majani 2 hadi 3 mpaka mwisho wa "*tillering*").

NB: Dalili huweza kuonekana baada ya siku 3 hadi 10, na baada ya wiki 2 hivi kwekwe hufa. Epuka kuzidisha kunakosababishwa na mwenendo wa polepole, kugeuka, kusimama ama kurudia.

KUTAYARISHA DAWA

Jaza nusu tanki la kunyunyizia na maji. Ongeza kipimo kinachohitajiwa cha GLEAN® 75 DF kwenye hilo tanki la kunyunyizia huku ukitingiza kwa nguvu. Ongeza mpaka kipimo kinachohitajika kwenye tanki la kunyunyizia.

KUSAFISHA KINYUNYIZIO

Ni muhimu kusafisha kabisa kila tone la GLEAN® 75 DF mara baada ya matumizi ya vifaa hivyo. Kwanza safisha tanki, puliza tanki na nusu lita ya

chlorine bleach (5% sodium hypochlorite) ichanganywe na lita 100 za maji. Kwa muda wa dakika 15 weka kando mchanganyiko huu kabla ya kuutumia. kisha baada ya kuondoa vifuniko vya kinyunyizio vioshe tena kila kimoja kwa uangalifu. Kumaliza, safisha tanki pamoja na vichungi kwa maji ya bomba ili kuweza kutoa kila tone lililobaki la chlorine bleach.

MIMEA INAYOFUATIA

Wakati wa kufuatia mimea, majarabio kuendelea, inapendekezwa kwamba ngano ama shayiri ipandwe mara moja kufuatia mimea ya nafaka iliyonyunyiziwa dawa ya GLEAN® 75 DF. Kama mimea haikuota kwa sababu yoyote ile basi shamba inabidi lipandwe tena na ngano au shayiri pekee.

AINA YA KWEKWE

Majarabio Kenya yameonyesha kwamba aina ya kwekwe wafuatao wanazuliwa na GLEAN® 75 DF.

Common Name

Oxalis

Spurrey

Chickweed

Wild raddish (white charlock)

Scientific Name*Oxalis spp**Spergula arvensis**Stellaria media**Raphanus raphanistrum*

Black bindweed
 Round-leaved Goosefoot
 Prostrate knotweed
 Shepherd's purse
 Pigweed
 Mallow
 Chinese lantern
 Mexican marigold
 Gallant soldier
 Devils thorn
 Black knightshade
 Thorn apple
 Blackjack
 Portulaca
 Volunteer Rape Double thorn
 Rye grass

Polygonum Convolvulus
Chenopodium opulifolium
Polygonum aviculare
Capsella bursa-pastoris
Amaranthus spp
Malva verticillata
Nicandra physalodes
Tagetes minuta
Galinsoga parviflora
Emex australis
Solanum nigrum
Datura stramonium
Bidens pilosa
Portulaca oleracae
Oxygonum sinuatum
Lolium spp.

TAHADHARI

Fungia mbali na watoto. Dawa hii inaweza kuwasha macho, pua, koo na ngozi. Chukua tahadhari wakati wakunyunyizia dawa hii. Manyunyu yake yasiangukie mimea mingine hasa yenye majani mapana.

Baada ya kutumia dawa hii: Osha *gloves* vizuri na kisha unawe mikono kwa sabuni na maji kisha ubadilishe nguo.

KUHIFADHI

Hifadhi ndani ya chombo chake halisi pahali pasipo na joto na pakavu mbali na watoto, vyakula vya binadamu na wanyama. Usiweke karibu na chakula, vinywaji, au chakula cha wanyama.

UTUPAJI

Usitumie chombo hiki tena. Zika au choma pahali penye panaruhusiwa.

MADHARA KWA MAZINGARA

Usimwage mabaki ya dawa hii, au kusunza chombo kilicho kuwa na dawa hii, ndani ya mito, visima au vidimbwi. Epusha ukungu kwa mazao, mahali pa kulisha mifugo au sehemu zilizo na chemi chemi za maji.

MAELEZO YA HUDUMA YA KWANZA

Ikimwagikia Ngozi: Vua nguo na viatu vilivyo mwagikiwa na dawa mara moja. Osha vizuri kwa kutumia maji ya kutosha na sabuni. Pata ushauri wa daktari .
Ikiingia kwa macho: Yasafishe macho kwa maji yanayo tiririka kwa muda wa dakika 10 huku ukiyaacha wazi. Muone daktari .



Ikivutwa: Anza kumpumulisha mgojwa kwa njia zilizopendekezwa na uedelee mpaka daktari aanze kumhudumia.

Ikimezwa: Mtapishe kwa kutia kidole ndani ya koo kwa utaratibu. Rudia mpaka matapiko yaishe dawa. Pata ushauri wa daktari.

MAELEZO YA SUMU /ARIFA KWA DAKTARI

Makata: Hakuna makata inayojulikana. Tibu mjeruhi kulingana na dalili.

Wakati wa dharura piga simu bila malipo 0800 730030/0800 720021 (masaa 24).

ILANI KWA MTUMIAJI

Dawa hii lazima itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyu salama.

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokea kutokana na matumizi ya dawa hii, kwenye binadamu au mali na lazima aikubali dawa hii katika hali hiyo.





NOIR



