

K-36637/31505 - TANZANIA - PAGE 2 (BASE)

DESCRIPTION

AVAUNT® 150 EC is an insecticide with stomach and contact action mainly against larvae of Bollworm (*Helicoverpa armigera*) and Army worms (*Spodoptera* spp) in tomatoes, beans, cucurbits, cabbage crops and potatoes, and diamond back moth larvae (*Plutella xylostella*) on cabbage crops and Bud worms (*Helicoverpa armigera*) in tobacco.

® Registered trademark of DuPont



**Emulsifiable concentrate
insecticide containing 150 g/L Indoxacarb**

Registration no: IN/0385
(Approved by TPRI)
(Imbethibitiswa na TPRI)



MAELEZO YA JUMLA

AVAUNT® 150 EC ni kiuadudu chenye kuwa kwa kumezwa au kugusa mwili wa viwavi vya Bollworm, viwavi jeshi katika mazao ya nyanya, tumbaku, maharage, matangi, kabichi na viazi mwingiro na viwavi vya diamond back moth katika kabichi na Budworms (*Helicoverpa armigera*) kwenye tumbaku.

Distributed by :

Tanzania Crop Care Ltd
Njiro, Arusha Tanzania
Phone: +255 27 2549054
Fax: +255 27 2549066

Produced by / Imetengenezwa na : Du Pont - USA
Validity / Maisha rafuni : 2 years from manufacturing date / Miaka 2 baada ya tarehe ya kutengenezwa.

K-36637/31505 - TANZANIA



HARMFUL

CYAN MAGENTA JAUNE NOIR





K-36637/31505 - TANZANIA - PAGE 3

Emulsifiable concentrate insecticide containing 150 g/L Indoxacarb

Registration no: IN/0385 (Approved by TPRI)

RATES OF APPLICATION

AVAUNT® 150 EC is applied post-emergence to crops, using standard equipment, 1 to 4 times with a 7-10 day interval depending on the crop and its growth stage. The recommended rate is 0.25 L (formulated product)/Ha i.e. 37.5 g Indoxacarb (active ingredient)/Ha.

As long as an application of **AVAUNT® 150 EC** has dried on the plant there is no need to re-treat in the event of rain.

PRE HARVEST INTERVAL

- Tomatoes, Beans, Cucurbits & Cabbage Crops: 3 days before harvest.
(for local consumption only. For crops destined for export contact your local DuPont representative for advise before applying **AVAUNT® 150 EC**).
- Tobacco: 28 days before harvest.

NOIR





K-36637/31505 - TANZANIA - PAGE 4

MIXING INSTRUCTIONS:

Start with a clean sprayer, half fill the spray tank with clean water. Thoroughly shake the container of **AVAUNT® 150 EC** to ensure an uniform suspension, add the required amount of the product and start agitation. Complete the filling of the spray tank with water. Continue agitation during spraying. Do not prepare more spray liquid than necessary.

COMPATIBILITY:

AVAUNT® 150 EC is compatible with a range of insecticides, herbicides, and fungicides used in all the above listed crops.

Please refer to DuPont's National distributor Tanzanian Crop Care Limited for the latest information on supported compatibilities.

SAFETY PRECAUTIONS

- Harmful if swallowed: Keep away from foodstuff and water supplies.
- Always wear protective clothing and rubber gloves while handling **AVAUNT® 150 EC**.
- Do not eat, drink or smoke in working area or in the chemical store.
- Avoid contact with eyes, skin, or clothing.

NOIR





- Avoid inhaling spray mist or vapors.
- Wash thoroughly with soap and water after handling before eating or smoking.
- Treated areas can be re-entered with protective clothing as soon as the product has dried on the leaves.
- Special care must be taken to avoid spray drifts outside the target area, or onto ponds, waterways or ditches.
- Immediately after application, completely drain spray equipment, thoroughly rinse sprayer and flush the hoses, boom and nozzles with clean water. Take all necessary safety precautions when cleaning equipment; do not clean near wells, water sources.

FIRST AID INSTRUCTIONS

- Eye contact: Wash with plenty of water for 15 to 20 minutes, and then get medical attention.
- If inhaled and breathing has stopped: Perform artificial respiration, and remove from area immediately. Keep the person warm and at rest. Get medical attention.
- If ingested: Seek medical attention immediately.
- Skin contact: Wash off immediately with soap and plenty of water. Take off contaminated clothing and shoes immediately. In the case of skin irritation or allergic reactions see a physician. Wash contaminated clothing before re-use.

NOIR





K-36637/31505 - TANZANIA - PAGE 6

STORAGE

Store safely in under lock and key in a cool, dry, well ventilated place away from children, animals, food, feedstuffs and drinking water.

DISPOSAL INSTRUCTIONS

Do not use empty containers for any other purpose. Triple rinse empty containers and dispose of safely according to local regulations.

RESISTANCE MANAGEMENT:

Some pests are known to develop resistance to products used repeatedly for control. When this occurs, the recommended dosages fail to suppress the pest population below the economic threshold. The development of resistance cannot be predicted and local advisers should be consulted for detailed recommendations. In general the following points will help to avoid resistance:

1. Alternate compounds of different chemical classes.
2. Monitor pest populations and apply **AVAUNT® 150 EC** when locally determined economic thresholds are reached.
3. Follow label recommendations for rates and applications intervals.

NOIR





K-36637/31505 - TANZANIA - PAGE 7

NOTICE TO THE BUYER

Manufacturer warrants that the product conforms to the chemical description on the label when it is presented in tightly closed original packaging. We are not liable for direct or indirect consequences resulting from improper storage or use contrary to regulations. This pest control product is to be used only in accordance with the directions on this label.



NOIR





K-36637/31505 - TANZANIA - PAGE 8

Emulsifiable concentrate insecticide containing 150 g/L Indoxacarb

Registration no: IN/0385 (Imbethibitiswa na TPRI)

KIWANGO CHA KUTUMIA

Tumia **AVAUNT® 150 EC** lita 0.25 kwa hekari yaani gramu 37.5 Indoxacarb/hekari mara 1-4 kila baada ya siku 7-10 kutegemea aina ya zao na hali halisi ya mashambulizi ya wadudu.

Ikiwa **AVAUNT® 150 EC** imekaukia kwenye mimea usipulizie tena wakati wa kunyesha mvua.

MUDA KABLA YA KUVUNA

Nyanya, Maharage, Matango, Kabichi	siku 3 kabla ya kuvuna
Tumbaku	siku 28 kabla ya kuvuna

JINSI YA KUCHANGANYA DAWA.

Tikisa chupa ya dawa vizuri kabla ya kuweka dawa kwenye bomba la kupulizia.

NOIR





Weka maji safi kwenye bomba la kupulizia. Weka kiasi kinachohitajika cha **AVAUNT® 150 EC** kwenye bomba la kupulizia, koroga vizuri na ongeza kiasi cha maji kinachohitajika. Endelea kukoroga unapochanganya au kupuliza dawa hadi dawa iishe kwenye bomba.

Tumia bomba safi ya kupulizia, weka maji safi nusu tank. Vizuri tikisa chupa ya AVAUNT® 150 EC na uweke kiasi cha dawa kinachohitajika katika tank yenye maji na tikisa tank vizuri kupata mchanganyiko wa dawa na maji. Anza kupuliza kwenye mimea.

ULINGANIFU/UPATANIFU

AVAUNT® 150 EC inawezwa kuchanganywa na viudadu, viuagugu, viuakuvu vyengine vinavyotumika katika mazao yaliyotajwa hapo juu. Usichanganye **AVAUNT® 150 EC** an madawa mengine yenye alikali (kama magadi, potashi).

TAHADHARI

- Vaa mavazi ya kinga wakati wa kupiga dawa.
- Usile, usinywe wala kuvuta sigara wakati wa kupulizia dawa.
- Jihadhari dawa isiguse macho, ngozi, nguo.
- Jihadhari usivute mvuke wa dawa.

NOIR





K-36637/31505 - TANZANIA - PAGE 10

- Oga baada ya kupulizia dawa, kabla ya kula au kuvuta sigara.
- Usioshe vyombo vyenye dawa karibu na chanzo cha maji kama kisima, maziwa, mito.

HUDUMA YA KWANZA

Ikiingia machoni: Osha kwa sabuni na maji mengi, kisha muone daktari.

Akivuta mvuke au vitone vya dawa: Muondoe mgonjwa na mpeleke sehemu yenye hewa safi, kisha muone daktari.

Ikimezwa: Muone daktari

Kikata sumu: Atropine na mchanganyiko wa Oximes

Ikigusa kwenye ngozi: Osha kwa sabuni na maji mengi

KUHIFADHI

Hifadhi kwenye chombo chake asili. Fungia kwa kufuli mbali na watoto, chakula, maji na vinywaji, penye hewa safi na pakavu

TAARIFA KWA MTUMIAJI

Dawa hii lazima itumiwe kufuatana na maelezo yaliyo andikwa juu ya kibandiko.

NOIR





K-36637/31505 - TANZANIA - PAGE 11 (REAR COVER)



NOIR






K-36637/31505 - TANZANIA - PAGE 12 (COVER)

OPEN HERE

DESCRIPTION

AVAUNT® 150 EC is an insecticide with stomach and contact action mainly against larvae of Bollworm (*Helicoverpa armigera*) and Army worms (*Spodoptera* spp) in tomatoes, beans, cucurbits, cabbage crops and potatoes, and diamond back moth larvae (*Plutella xylostella*) on cabbage crops and Bud worms (*Helicoverpa armigera*) in tobacco.

® Registered trademark of DuPont



HARMFUL



**Emulsifiable concentrate
insecticide containing 150 g/L Indoxacarb**

100 ml

Registration no: IN/0385
(Approved by TPRI)
(Imbethibitiswa na TPRI)

MAELEZO YA JUMLA

AVAUNT® 150 EC ni kiuadudu chenye kuua kwa kumezwa au ku-gusa mwili wa viwavi vya Bollworm, viwavi jeshi katika mazao ya nyanya, tumbaku, maharage, matango, kabichi na viazi miringo na viwavi vya diamond back moth katika kabichi na Budworms (*Helicoverpa armigera*) kwenye tumbaku.

Distributed by : Tanzania Crop Care Ltd

Njiro, Arusha Tanzania

Phone: +255 27 2549054

Fax: +255 27 2549056

Produced by / Imetengenezwa na : Du Pont - USA

Validity / Maisha rafuni : 2 years from manufacturing date

/ Miaka 2 baada ya tarehe ya kutengenezwa.

K-36637/31505 - TANZANIA



Lot nr.:



CYAN MAGENTA JAUNE NOIR

